

GENERAL MATHEMATICS 11

Name of Learner: _____ Grade Level: _____

Section: _____ Date: _____

LEARNING ACTIVITY SHEET

EVALUATES A FUNCTION

Background Information for the Learners

In this topic, you will learn about evaluating functions. To evaluate a function is simply to *substitute/replace* a certain variable with a given number or expression, in this case we refer this as the x variable. We then write $f(a)$ to show that f is being evaluated at a for some a in the domain of f .

Example 1.

Evaluate the given functions at $x = 2$

1. $f(x) = 5x + 8$
2. $g(x) = x^2 + 3x - 4$
3. $h(x) = |x + 3| - 2$
4. $t(x) = x^3 - 5x^2$
5. $p(x) = [x] + 5$, where $[x]$ is the smallest integer greater than or equal x .

Solution:

1. $f(2) = 5(2) + 8 = 10 + 8 = 18$
2. $g(2) = (2)^2 + 3(2) - 4 = 4 + 6 - 4 = 6$
3. $h(2) = |2 + 3| - 2 = |5| - 2 = 5 - 2 = 3$
4. $t(2) = (2)^3 - 5(2)^2 = 8 - 20 = -12$
5. $p(2) = [2] + 5 = 2 + 5 = 7$

Learning Competency

Evaluates a Function (GM_M11GM-Ia-2)

EXERCISE 1

Directions: Evaluate the following functions at $x = 3.5$.

[1 point each]

1. $f(x) = 7x - 1$
2. $g(x) = x^2 - 4$
3. $h(x) = |4x - 21| - 2$
4. $t(x) = 2 \cdot 3^{x-1.5}$
5. $s(x) = \sqrt{x^2 - 2x + 4}$
6. $r(x) = -2^{-2x}$
7. $w(x) = 3^{2x-5}$
8. $q(x) = \lfloor x \rfloor + 4$
9. $f(x) = \frac{x+4}{x^2-2}$
10. $f(x) = \begin{cases} 5x & , x < 2 \\ x^2 - 1 & , 2 \leq x < 5 \\ x^3 + 2 & , x \geq 5 \end{cases}$

EXERCISE 2

Directions: Evaluate each function.

[1 point each]

1. $f(x) = |x + 2| + 4$, find $f(-8)$
2. $h(x) = 3^{3x-2}$, find $h(2)$
3. $q(x) = 4^{3x}$, find $q(\frac{1}{2})$

For numbers 4-5, use the function below

$$f(x) = x^2 + 5x + 4$$

4. Find $f(2)$
5. Find $f(x + 2)$

For numbers 6-8, use the function below

$$f(x) = \begin{cases} x + 2 & , x \leq -4 \\ \sqrt{16 - x^2} & , -4 < x < 4 \\ 2 - x & , x \geq 4 \end{cases}$$

6. Find $f(-5)$
7. Find $f(2)$
8. Find $f(5)$

For items 9-10, given $f(x) = \sqrt{x+4}$,

9. Find $f(x^2 - 4)$

10. Find $\frac{f(x+h)-f(x)}{h}, h \neq 0$

EXERCISE 3

Directions: Solve the given problems

[3 points each]

1. An internet café charges P25.00 per hour (or a fraction of an hour) for the first four hours and an extra of P15.00 per hour for each next hour. What is the amount paid for following time?
 - a. 2 hrs.
 - b. 3.5 hrs.
 - c. 6 hrs.
 - d. 7.2 hrs.
2. In a certain city, the Philippine Statistics Office has recorded a total population of 680,000 in the year 2018. The population increases at the rate of 0.25% annually. Suppose the rate of increase is constant. Determine the population in the following years.
 - a. 10 yrs.
 - b. 14 yrs.
 - c. 17 yrs.
3. Driving lessons require a rental car fee of P 500.00 for the first 8 km. and for every kilometer added charges an additional fee of P 50.00. How much will it cost for the following distance.
 - a. 7 km.
 - b. 9 km.
4. A stone is thrown straight up from a building with a velocity, in m/s, given by $V(t) = 15 - 9.8t$. After t seconds, calculate the following velocities. Make an interpretation of the results.
 - a. $V(0)$
 - b. $V(1)$
5. In particular bacteria culture, if $p(x)$ bacteria are present at x minutes, then $p(x) = Ae^{0.03x}$ such that A is a constant. Initially, there are 1200 bacteria present, calculate the number of bacteria after:
 - a. 30 minutes
 - b. 1 hr.
 - c. 1.5 hrs.

Reflection:

What have you learned from this topic?

References:

Verzosa, D.B, et.al (2016). General Mathematics. Quezon City, Manila
Alferez, G. S. (2014). Introduction to Calculus. Quezon City, Manila
Leithold, L. (1996). The Calculus 7. New York City.
<https://cdn.kutasoftware.com/Worksheets/Alg2/Evaluating%20Functions.pdf>

Answer Key

Exercise 1

- | | |
|---------|---------------------|
| 1. 23.5 | 6. $-\frac{1}{128}$ |
| 2. 8.25 | 7. 9 |
| 3. 5 | 8. 7 |
| 4. 18 | 9. 0.73 |
| 5. 9.25 | 10. 11.25 |

Exercise 2

- | | |
|--------------------|---|
| 1. 10 | 6. -3 |
| 2. 8 | 7. $2\sqrt{3}$ |
| 3. 8 | 8. -3 |
| 4. 18 | 9. x |
| 5. $x^2 + 9x + 18$ | 10. $\frac{1}{\sqrt{x+h+4} + \sqrt{x+4}}$ |

Exercise 3

- P50.00
 - P100.00
 - P130.00
 - P160.00
- $P = 697, 193$
 - $P = 704, 191$
 - $P = 709, 485$
- P500.00
 - P550.00
- 15 m/s
 - 5.2 m/s

This implies that the initial velocity is 15 m/s and gets more slowly at 5.2 m/s.
- $P(x) = 2, 952$
 - $P(x) = 7, 260$
 - $P(x) = 17, 856$